



2X SUPERSTRONG PCR MASTER MIX

User Guide

Catalog Number	Pack Size
ZT-SPCRMX-1	1,25 mL (100RXN)
ZT-SPCRMX-5	5 X 1,25 mL (5x100RXN)

Content and Storage Condition

Content	Quantity	Transportation Condition	Storage Condition
2X SUPERSTRONG PCR MASTER MIX	1,25 mL	-20°C	-20°C

Features

Concentration: 2X

Application: Molecular Biology

Purity Class: Molecular Biology

Appearance: Colorless, clear, liquid

Classification: General Substance

Reaction Speed: Standard

Polymerase: Taq Polymerase

2X PCR Master Mix Composition

0.05 U/ μ L Taq DNA polymerase, 10X Reaction buffer, 4 mM MgCl₂, 0.4mM each dATP, dCTP, dGTP and dTTP
It is a ready-to-use mixture.

Its half-life is more than 40 minutes at 95 ° C.

Produces 3'-dA PCR products

Contains modified nucleotides (for example, biotin-, digoxigenin-, fluorescently labeled nucleotides)

Applications

Routine PCR amplification of DNA fragments up to 5 kb in length and high throughput PCR.



Product Description

SOFTEC 2X SUPERSTRONG PCR Master Mix is a 2X concentration mixture consisting of all the components required for PCR except DNA template and primers. This pre-mixed formulation saves time and reduces contamination due to the reduction of pipetting steps required for PCR setup. The mixture is optimized for efficient and reproducible PCR. It is thermostable. Its half-life is more than 40 minutes at 95 °C.

Note

The error rate of Taq DNA Polymerase in PCR is 2.2×10^{-5} errors per nucleotide in each cycle. The accuracy of PCR is 4.5×10^4 . Accuracy is the inverse of the error rate and refers to the average number of correct nucleotides included before an error occurs.

Protocol

1. Gently vortex and briefly centrifuge SOFTEC 2X SUPERSTRONG PCR Master Mix after thawing on ice.
2. Place your 0.2ml microcentrifuge tubes on the ice and follow the table below for each 25 µL reaction volume.

2X SUPERSTRONG PCR	12,5 µL
Forward Primer	0,5 µL (10µM)
Reverse Primer	0,5 µL (10µM)
Template DNA	10pg - 1µg
Nuclease free water	to 25µL

3. Gently vortex the samples and spin down.
4. When using a thermal cycler that does not contain a heated lid, overlay the reaction mixture with 25 µL of mineral oil.
5. Perform PCR by following the table below for recommended thermal cycling conditions:

Step	Temperature (°C)	Time	Number of cycle
Pre-denaturation	95	1-3 min	1
Denaturation	95	30 s	25-40
Annealing	Tm-5	30 s	
Extension	72	1 min /1kb	
Final extension	72	5-15 min	1

Recommendations for Preventing Contamination for the PCR Reaction

- Perform the preparation of the DNA sample, preparation of the PCR mixture, use of the thermal cycling device and analysis operations in separate areas.
- Perform the preparation of the PCR mixture in a laminar flow cabinet with a UV lamp.
- Wear clean laboratory gloves and renew your gloves in different steps.
- Always perform “no template control” (NTC) reactions to check for contamination.

Tips

For GC-rich DNA templates, DNA denaturation time of 30 seconds can be prolonged to 3–4 min. Primer annealing temperature should be 5°C lower than the melting temperature (T_m) of the primers.

Annealing for 30 seconds is normally sufficient. If non-specific PCR products appear, the annealing temperature should be optimized stepwise in 1–2°C increments.

The recommended extension step is 1 min/kb at 72° for PCR products up to 2 kb. For larger products, the extension time should be prolonged by 1 min/kb.

For less than 10 copies of the template in the reaction, 40 cycles are recommended.

For higher template amounts, 25–35 cycles are sufficient.

Warning

Do not take delivery of the product whose packaging is damaged and do not use it.

CERTIFICATE OF ANALYSIS

Endodeoxyribonuclease Assay: No detectable degradation of DNA was observed after incubation of 1 µg of pUC19 DNA with 25 µL of SOFTEC 2X SUPERSTRONG PCR Master Mix in 50 µL of reaction mixture for 4 hours at 37°C and at 70°C.

Exodeoxyribonuclease Assay: No detectable degradation of DNA was observed after incubation of 1 µg of lambda DNA/HindIII fragments with 25 µL of SOFTEC 2X SUPERSTRONG PCR Master Mix in 50 µL of reaction mixture for 4 hours at 37°C and at 70°C.

Ribonuclease Assay: No contaminating RNase activity was detected after incubation of 1 µg of [3H]-RNA with 25 µL SOFTEC 2X SUPERSTRONG PCR Master Mix in 50 µL of reaction mixture for 4 hours at 37°C and at 70°C.

Quality authorized by: Zet Medical R&D Lab